arXiv:1511.03716v1 [math.GM] 11 Nov 2015

Evaluations of certain theta functions in Ramanujan theory of alternative modular bases

N.D. Bagis
Aristotele University of Thessaloniki
Thessaloniki Greece
nikosbagis@hotmail.gr

abstract

We give evaluations of certain Borwein's theta functions which appear in Ramanujan theory of alternative elliptic modular bases. Most of this theory where developed by B.C. Berndt, S. Bhargava and F.G. Garvan. We also study the most general class of these theta functions and give evaluation conjectures.

keywords: Ramanujan; theta functions; alternative modular bases; Evaluations;

1 Introduction

The Borwein-Borwein theorem for the cubic modular base ${}_2F_1\left(\frac{1}{3},\frac{2}{3};1;x\right)$, states that (see [6]):

Theorem 1. Let $\omega=\exp(2\pi i/3)$, $q=\exp(-\pi\sqrt{r})$, $r>0$ and

$$a_0(q):=\sum_{m,n=-\infty}^{\infty} q^{m^2+mn+n^2}, \tag{1}$$

$$b_0(q):=\sum_{m,n=-\infty}^{\infty} \omega^{m-n} q^{m^2+mn+n^2}, \tag{2}$$

and

$$c_0(q):=\sum_{m,n=-\infty}^{\infty} q^{(m+1/3)^2+(m+1/3)(n+1/3)+(n+1/3)^2}. \tag{3}$$

Then the solution of equation

$$\frac{{}_2F_1\left(\frac{1}{3},\frac{2}{3};1,1-x\right)}{{}_2F_1\left(\frac{1}{3},\frac{2}{3};1,x\right)}=\sqrt{r} \tag{4}$$

is

$$x = \alpha_r = \left(\frac{c_0\left(q^{2/\sqrt{3}}\right)}{a_0\left(q^{2/\sqrt{3}}\right)} \right)^3 \tag{5}$$

and

$$a_0^3(q) = b_0^3(q) + c_0^3(q) \tag{6}$$

Also in view of [6] which the case of Ramanujan elliptic modular bases theory is treated extesively, many results can be extracted. In this work we explore some of these ideas. From our part we give formulas for closed form evaluations of the above theorem theta functions.

2 Some Results

It is known from [6] p.82, the folowing relation of hypergeometric functions

$$\sqrt[4]{1+8x} \cdot {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; 1, x\right) = {}_2F_1\left(\frac{1}{6}, \frac{5}{6}; 1; \frac{1}{2} - \frac{1-20x-8x^2}{2(1+8x)^{3/2}}\right). \tag{7}$$

Seting $x = \alpha_{3r}$ in (7) and using (see [2] Theorem 4)

$$\beta_r = \frac{1}{2} - \frac{1-20\alpha_{3r}-8\alpha_{3r}^2}{2(1+8\alpha_{3r})^{3/2}}, \tag{8}$$

we get

$$\sqrt[4]{1+8\alpha_{3r}} \cdot {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; 1; \alpha_{3r}\right) = {}_2F_1\left(\frac{1}{6}, \frac{5}{6}; 1; \beta_r\right). \tag{9}$$

Using the identities $\alpha_{1/r} = \alpha'_r$ and $\beta_{1/r} = \beta'_r$ one can easily see that the multiplier of 3rd-degree in the 6-base (signature 6) is

Theorem 2.

$$m_3(6) = \frac{{}_2F_1\left(\frac{1}{6}, \frac{5}{6}; 1; \beta_{9r}\right)}{{}_2F_1\left(\frac{1}{6}, \frac{5}{6}; 1; \beta_r\right)} = \sqrt{3r} \cdot \sqrt[4]{\frac{1+8\alpha'_{3r}}{1+8\alpha_{3r}}} \tag{10}$$

From [6] Theorem 2.10 we have

Theorem 3. If $q = \exp(-\pi\sqrt{r})$, $r > 0$ and $z_r = {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; 1; \alpha_r\right)$, then

$$z_{3r} = a_0(q^2). \tag{11}$$

From [6] we also have

$$28(b_0^6(q^2) - c_0^6(q^2)) = 27N(q^6) + N(q^2), \tag{12}$$

where $N(q)$ is the Eisenstein series

$$N(q) = 1 - 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1-q^n}. \tag{13}$$

Also

$$N(q^2) = (1 - 20\alpha_{3r} - 8\alpha_{3r}^2) z_{3r}^6, \tag{14}$$

$$N(q^6) = \left(1 - \frac{4}{3}\alpha_{3r} + \frac{8}{27}\alpha_{3r}^2\right) z_{3r}^6 \tag{15}$$

and

$$b_0^6(q^2) - c_0^6(q^2) = (1 - 2\alpha_{3r}) z_{3r}^6. \tag{16}$$

Theorem 4. The number of rerpresentations $s(n)$ of the integer n in the form

$$n = x^2 + y^2 + xy + zw + z^2 + w^2 \tag{17}$$

is

$$s(n) = 12 \sum_{d|n} d - 12 \sum_{d|3, d|n} d \tag{18}$$

and $s(0) = 1$.

Proof. In [5], p.460, Entry 3(i) it has shown that

$$a_0(q)^2 = 1 + 12 \sum_{n=1}^{\infty} \frac{nq^n}{1-q^n} - 36 \sum_{n=1}^{\infty} \frac{nq^{3n}}{1-q^{3n}} \tag{19}$$

Using (1) along with the formula

$$\sum_{n=1}^{\infty} \frac{X(n) q^n}{1-q^n} = \sum_{n=1}^{\infty} \left(\sum_{d|n} X(d) \right) q^n, \tag{20}$$

we get the result.

Theorem 5. The number of representations of the integer n in the form

$$n = x^2 + xy + y^2 \tag{21}$$

is

$$r(n) = 6 \sum_{d \equiv 1(3), d|n} 1 - 6 \sum_{d \equiv 2(3), d|n} 1 \tag{22}$$

and $r(0) = 1$.

Proof. From the proof of [6] Theorem 2.12 p.11-12 and (11),(1) of the present article, we have

$$\sum_{m,n=-\infty}^{\infty} q^{m^2+mn+m^2} = 1 + 6\sum_{n=0}^{\infty} \frac{q^{3n+1}}{1-q^{3n+1}} - 6\sum_{n=0}^{\infty} \frac{q^{3n+2}}{1-q^{3n+2}}. \qquad (23)$$

Expanding the above formula into power series using (20) we get the result.

For the 5th base we have the relation

Theorem 6. If $q = e^{-\pi\sqrt{r}}$, $r > 0$ we have

$$u(\beta_r)^4 = 1 + 240\sum_{n=0}^{\infty} \frac{n^3 q^{2n}}{1-q^{2n}} = \frac{1}{2}\left(a^8 + b^8 + c^8\right) \qquad (24)$$

where $a = \theta_2(q) = \sum_{n=-\infty}^{\infty} q^{(n+1/2)^2}$, $b = \theta_3(q) = \sum_{n=-\infty}^{\infty} q^{n^2}$, and $c = \theta_4(q) = \sum_{n=-\infty}^{\infty}(-1)^n q^{n^2}$, $|q| < 1$.

From (20) and [3],[4] we also have

$$a_0(q)^2 = 1 + 12\sum_{n=1}^{\infty} \frac{nq^n}{1-q^n} - 36\sum_{n=1}^{\infty} \frac{nq^{3n}}{1-q^{3n}} =$$

$$= 1 - 12q\frac{d}{dq}\log\left(\eta(q^3)^{-1}\vartheta\left(\frac{3}{2},\frac{1}{2};q\right)\right) \qquad (25)$$

where

$$\vartheta\left(\frac{3}{2},\frac{1}{2};q\right) = q^{1/12}\eta(q^3)Q(k_r^2)^{1/12} \qquad (26)$$

The minimal polynomial of $u = Q(v)$ is

$$-16u^3 + 387420489v + 19131876uv + 196830u^2v + 84u^3v + u^4v - 1549681956v^2 -$$

$$-76527504uv^2 - 787320u^2v^2 - 12480u^3v^2 - 4u^4v^2 + 2324522934v^3 + 114791256uv^3 +$$

$$+1180980u^2v^3 - 40712u^3v^3 + 6u^4v^3 - 1549681956v^4 - 76527504uv^4 - 787320u^2v^4 -$$

$$-12480u^3v^4 - 4u^4v^4 + 387420489v^5 + 19131876uv^5 + 196830u^2v^5 +$$

$$+ 84u^3v^5 + u^4v^5 - 16u^3v^6 = 0 \qquad (27)$$

and is solvable. Hence

$$a_0(q)^2 = -q\frac{d}{dq}\log Q(k_r^2) \qquad (28)$$

hence

$$a_0(q)^2 = -q\frac{d}{dk}\left(\log Q(k^2)\right)\frac{dk}{dq} \qquad (29)$$

But from [5] chapter 17 Entry 9 p.120 we have

$$\frac{dq}{dk} = \frac{q\pi^2}{2k(1-k^2)K(k)^2}. \tag{30}$$

Setting (here the prime on Q means derivative):

$$\lambda_r := \sqrt{-\frac{Q'(k_r^2)}{Q(k_r^2)}} \tag{31}$$

we get the next

Theorem 7. If $u = Q(v)$ is that of (26) and $q = \exp(-\pi\sqrt{r})$, then

$$a_0(q) = \sum_{n,m=-\infty}^{\infty} q^{m^2+mn+n^2} = \frac{2K}{\pi} k_r k_r' \lambda_r \tag{32}$$

Also

$$a_0(q^2) = \sum_{n,m=-\infty}^{\infty} q^{2[m^2+mn+n^2]} = \frac{2K}{\pi} \sqrt[4]{\frac{1-k_r^2+k_r^4}{1+8\alpha_{3r}}} \tag{33}$$

Moreover if $r-$positive rational, then

$$\frac{\pi^2 a_0(q)^2}{K^2} = Algebraic \tag{34}$$

where $k' = \sqrt{1-k^2}$ is the complementary modulus and $Q'(x)$ is the first derivative of $Q(x)$.

Note. Relation (33) is known from [2] and (11),(5).

Example. Let $q = e^{-\pi}$, then $k_1 = \frac{1}{\sqrt{2}}$. Solving equation (26) (which is quartic) and setting $v = 1/2$ we get

$$Q\left(\frac{1}{2}\right) = 81\left(885 + 511\sqrt{3} - 3\sqrt{174033 + 100478\sqrt{3}}\right)$$

and

$$Q'\left(\frac{1}{2}\right) = 162\left(5082 + 2934\sqrt{3} - \sqrt{51655599 + 29823374\sqrt{3}}\right)$$

Also

$$K(k_1) = K\left(\frac{1}{\sqrt{2}}\right) = \frac{8\pi^{3/2}}{\Gamma\left(-\frac{1}{4}\right)^2}$$

Hence

$$a_0\left(e^{-\pi}\right) = \frac{8\sqrt{2\pi\left(\sqrt{24+14\sqrt{3}}-3\right)}}{\sqrt[4]{3}\Gamma\left(-\frac{1}{4}\right)^2} \tag{35}$$

From relation (5) we get the next

Corollary. If $q=\exp(-\pi\sqrt{r})$ with $r>0$, then

$$c_0(q^2)=\sum_{m,n=-\infty}^{\infty} q^{2[(m+1/3)^2+(m+1/3)(n+1/3)+(n+1/3)^2]}=$$

$$=\frac{2K}{\pi}\frac{1-k_r'}{1+k_r'}\sqrt{k_r'}\alpha_{3r}^{1/3}\lambda_{4r} \tag{36}$$

Also

$$b_0(q^2)=\sum_{m,n=-\infty}^{\infty}\omega^{m-n}q^{2[m^2+mn+n^2]}=\frac{2K}{\pi}\frac{1-k_r'}{1+k_r'}\sqrt{k_r'}\lambda_{4r}\sqrt[3]{1-\alpha_{3r}} \tag{37}$$

and consequently

Theorem 8. For $r>0$, we have

$$\alpha_{3r}=\frac{1}{8}\left(-1+\frac{(1-k_r^2+k_r^4)(1+k_r')^4}{(1-k_r')^4k_r'^2\lambda_{4r}^4}\right) \tag{38}$$

Theorem 9. The equation

$$Cx(1-x)^2=1 \tag{39}$$

have solution

$$x=\frac{1}{2}\frac{\theta_3^8(q)}{\theta_2^8(q)+\theta_3^8(q)+\theta_4^8(q)} \tag{40}$$

where $C=\frac{j_r}{256}$, where j_r is the $j-$invariant.

Proof. It holds (see[1] p.204)

$$j_r=256\frac{\left(1-(k_rk_r')^2\right)^3}{(k_rk_r')^4} \tag{41}$$

Also from [theta-null]

$$d^8=a^8+b^8+c^8=2\left(\frac{2K}{\pi}\right)^4(1-(k_rk_r')^2) \tag{42}$$

Hence

$$k_rk_r'=\sqrt{1-\frac{d^8}{2\theta_3^8}} \tag{43}$$

Setting this value to (41) we get

$$j_r = 128\frac{d^{24}}{\theta_3^8\left(d^8 - 2\theta_3^8\right)^2}$$

But if $x = \frac{\theta_3^8}{2d^8}$, then

$$\frac{j_r}{256} = \frac{1}{(1-x)^2 x}$$

Hence we get the result.

From relation (9) we obtain solution to a quartic equation.

Theorem 10. The equation

$$t^4 + 8C_1 t^3 + 18t^2 - 27 = 0 \tag{44}$$

have solution

$$t = \sqrt{1 + 8\alpha_r} \tag{45}$$

where r is given from $C_1 = 1 - 2\beta_r$.

Proof. From (8) wew get that equation

$$\frac{1 - 20x - 8x^2}{(1+8x)^{3/2}} = C_1 \tag{46}$$

admits solution $x = \alpha_{3r}$ with r such that $C = 1 - 2\beta_r$.

3 Conjectures on Theta functions

Lemma. Assume the quadratic forms

$$Q_1 = a_1x^2 + b_1xy + c_1y^2 \text{ and } Q_2 = a_2x^2 + b_2xy + c_2y^2, \tag{47}$$

where a_i, b_i, c_i, integers with $a_i > 0$, $b_i \geq 0$, $c_i \geq 0$, for $i = 1, 2$.
If $b_1^2 - 4a_1c_1 = b_2^2 - 4a_2c_2 = -|D| < 0$, then Q_1 and Q_2 are equivalent, meaning that the number of representations of $n \in \mathbf{N}$ in Q_1 equals to that of Q_2.

Theorem 11. Suppose that a, b, c are non negative integers with $a \neq 0$ and $D = b^2 - 4ac = -|D| < 0$, then

$$\sum_{n,m=-\infty}^{\infty} q^{an^2+bnm+cm^2} = \frac{2K}{\pi}F\left(|D|, k_r\right),\ q = e^{-\pi\sqrt{r}},\ \forall r > 0 \tag{48}$$

where $F(x, y)$ is two variable algebraic function.

Theorem 12. Let a, b, c be non negative integers with $a \neq 0$. Set

$$a(q) := \sum_{n,m=-\infty}^{\infty} q^{an^2+bnm+cm^2},\ q = e^{-\pi\sqrt{r}},\ r > 0. \tag{49}$$

Then if $D = b^2 - 4ac = -|D| \in \{-3, -4, -7\}$, we have

$$a(q)^2 = \frac{1}{D+1}\left(1 - 24\sum_{n=1}^{\infty}\frac{nq^n}{1-q^n} + D\left(1 - 24\sum_{n=1}^{\infty}\frac{nq^{-Dn}}{1-q^{-Dn}}\right)\right). \tag{50}$$

Hence we get the following

Corollary. The number of representations of the integer $n \neq 0$ in the form

$$n = a(x^2 + z^2) + b(xy + zw) + c(y^2 + w^2), \tag{51}$$

with a, b, c non negative integers $a \neq 0$, $D = b^2 - 4ac = -|D| = -3, -4, -7$, is

$$s(n) = -\frac{24}{D+1}\sum_{d|n} d + \frac{24D}{(D+1)D}\sum_{d|n, -D|d} d. \tag{52}$$

Set now

$$k_i(x) := \left(\frac{K\left(\sqrt{1-x^2}\right)}{K(x)}\right)^2. \tag{53}$$

For all a, b, c non negative integers and $a \neq 0$, such that $D = -3$ obviously we have

$$\sum_{n,m=-\infty}^{\infty} q^{an^2+bnm+cm^2} = {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; 1; \alpha_{3r/4}\right). \tag{54}$$

Also for all $r = k_i\left(\frac{m}{n}\right)$, $m < n$, m, n positive integers, the minimal polynomial of

$$S_3 = S_3(x) = \frac{\pi}{2K(x)}\sum_{n,m=-\infty}^{\infty} q_1^{an^2+bnm+cm^2} = F(3, x), \tag{55}$$

$q_1 = e^{-\pi\sqrt{k_i(x)}}$, $x \in \mathbf{Q}$, $0 < x < 1$, has always degree equivalent to 4 and hence is solvable, since from Theorem 7 and (31) we have

$$S_3(x) = x\sqrt{1-x^2}\sqrt{-\frac{Q'\left(x^2\right)}{Q\left(x^2\right)}}. \tag{56}$$

Moreover

Theorem 13. One can experimentaly see that exists integer coefficients a_{mn}, c_n and $N_1, N \in \mathbf{N}$, such that

$$\sum_{m,n=0}^{N_1} a_{mn}\left(\sum_{l=0}^{N} c_l y^l\right)^m S_{|D|}(y)^n = 0,\ y \in \mathbf{R},\ 0 < y < 1, \tag{57}$$

where $S_{|D|}(x) = F(|D|, x)$, $D = b^2 - 4ac = -|D| < 0$, a, b, c non negative integers, $a \neq 0$.

Note to the reader that $A = \{x : x \in \mathbf{Q} \text{ and } 0 < x < 1\}$ is dense in $\mathbf{R}$.

Assume in general that a, b, c are integers with $a > 0$, $b \geq 0$, $c > 0$, $q = e^{-\pi\sqrt{r}}$, $r > 0$

$$\theta_3(q) = \sum_{n=-\infty}^{\infty} q^{n^2} = \sqrt{\frac{2K}{\pi}}. \tag{58}$$

Also set

$$u := \left(\frac{F\left(|D|, m(q)\right)}{\theta_3(q)}\right)^2 \text{ and } v := m(q) = k_r,\ r > 0, \tag{59}$$

then

1. For $D = -4$

$$u = 1 \tag{60}$$

2. For $D = -8$

$$-4u + 4u^2 + v = 0 \tag{61}$$

3. For $D = -12$

$$-1 - 8u - 18u^2 + 27u^4 + 16uv = 0 \tag{62}$$

4. For $D = -16$

$$-64u + 64u^2 - 256u^3 + 256u^4 + 48uv + 32u^2v + v^2 = 0 \tag{63}$$

5. For $D = -20$

$$-1+26u-275u^2+1500u^3-4375u^4+6250u^5-3125u^6-256uv+256uv^2 = 0 \tag{64}$$

6. For $D = -24$

$$-6912u^4 - 55296u^5 - 124416u^6 + 186624u^8 - 1024uv - 8192u^2v -$$
$$-18432u^3v + +6912u^4v + 69120u^5v + 62208u^6v + 1280uv^2 + 8480u^2v^2 +$$
$$+ 16128u^3v^2 + 4320u^4v^2 - 288uv^3 + 112u^2v^3 + v^4 = 0 \tag{65}$$

8. For $D = -28$

$$-1-48u-980u^2-10976u^3-72030u^4-268912u^5-470596u^6+823543u^8+2144uv-$$
$$-37632u^2v+21952u^3v+537824u^5v-6144uv^2+37632u^2v^2+4096uv^3 = 0 \tag{66}$$

9. For $D = -32$

$$-16384u + 16384u^2 - 589824u^3 + 589824u^4 - 6291456u^5 + 6291456u^6 -$$
$$-16777216u^7 + 16777216u^8 + 28672uv - 221184u^2v + 933888u^3v - 327680u^4v +$$
$$+4456448u^5v+1048576u^6v-13568uv^2+206336u^2v^2-372736u^3v^2+24576u^4v^2+$$
$$+1216uv^3 + 256u^2v^3 + v^4 = 0 \tag{67}$$

10. For $D=-36$

$$-16384u+16384u^2-589824u^3+589824u^4-6291456u^5+6291456u^6-$$
$$-16777216u^7+16777216u^8+28672uv-221184u^2v+933888u^3v-327680u^4v+$$
$$+4456448u^5v+1048576u^6v-13568uv^2+206336u^2v^2-372736u^3v^2+$$
$$+24576u^4v^2+1216uv^3+256u^2v^3+v^4=0 \tag{68}$$

11. For $D=-40$

$$12800000u^6-332800000u^7+3520000000u^8-19200000000u^9+56000000000u^{10}-$$
$$-80000000000u^{11}+40000000000u^{12}-262144uv+6815744u^2v-72089600u^3v+$$
$$+393216000u^4v-1153280000u^5v+1785600000u^6v-2163200000u^7v+$$
$$+6080000000u^8v-13600000000u^9v+12000000000u^{10}v+589824uv^2-$$
$$-13631488u^2v^2+126156800u^3v^2-588704000u^4v^2+1412480000u^5v^2-$$
$$-1489600000u^6v^2+355200000u^7v^2+540000000u^8v^2-425984uv^3+$$
$$+3899392u^2v^3-58668800u^3v^3+197984000u^4v^3-293280000u^5v^3+$$
$$+10400000u^6v^3+102400uv^4+2920752u^2v^4+4463200u^3v^4+102000u^4v^4-$$
$$-4200uv^5+504u^2v^5+v^6=0 \tag{69}$$

Theorem 14. If $D=-3,-7,-11,-19,-43,-67,-163$ we have

$$\sum_{n,m=-\infty}^{\infty} q^{an^2+bnm+cm^2}=1+p\sum_{n=1}^{\infty}\left(\frac{n}{-D}\right)\frac{q^n}{1-q^n},\ \forall|q|<1. \tag{70}$$

where $p=6,2,2,2,2,2$ respectively.

Corollary. Let a,b,c be non negative integers with $a\neq 0$. The number of representations of a positive integer n into forms

$$Q=ax^2+bxy+cy^2,\ x,y\in\mathbf{Z}, \tag{71}$$

with $D=-|D|=-3,-7,-11,-19,-43,-67,-163$ is

$$r(n)=p\sum_{d|n}\left(\frac{d}{-D}\right),\ \text{if } n\in\mathbf{N} \tag{72}$$

and $r(0)=1$ if $n=0$, where $p=6,2,2,2,2,2,2$ respectively.

References

[1]: J.V. Armitage W.F. Eberlein. 'Elliptic Functions'. Cambridge University Press. (2006)

[2]: N.D. Bagis, M.L. Glasser. 'Conjectures on the Evaluation of Alternative Modular Bases and Formulas Approximating $1/\pi$'. Journal of Number Theory. (Elsevier), (2012).

[3]: N.D. Bagis, M.L. Glasser. 'Conjectures on the evaluation of certain functions with algebraic properties'. Journal of Number Theory. 155 (2015), 63-84

[4]: N.D. Bagis. 'On the Complete Evaluation of Jacobi Theta Functions'. arXiv:1503.01141v1 [math.GM],(2015).

[5]: B.C. Berndt. 'Ramanujan's Notebooks Part III'. Springer Verlag, New York (1991).

[6]: B.C. Berndt, S. Bhargava, F.G. Garvan. 'Ramanujan's theories of elliptic functions to alternative bases'. Trans. Amer. Math. Soc. 347 (1995) 4163-4244.